

REPORT: NATIONAL EMISSIONS FROM L&G EQUIPMENT

Jamie L Banks, PhD, MS*

Quiet Communities, Inc., PO Box 533, Lincoln, MA 01773

Robert McConnell, Environmental Engineer

US Environmental Protection Agency, Region 1, 5 Post Office Square, Boston, MA 02109

ABSTRACT

Background: The contribution of gasoline-powered lawn and garden equipment (GLGE) to air pollutant emissions in the United States has not been extensively studied. **Goal:** Our goal is to provide annual US and state-level emissions estimates of volatile organic compounds (VOC): criteria pollutants (carbon monoxide [CO], nitrogen oxides [NO_x], particulate matter [PM] <10 microns, including PM < 2.5 microns [PM₁₀, PM_{2.5}]; and carbon dioxide (CO₂) from GLGE, with a focus on 2-stroke engines. **Methods:** Pollutant emissions data were extracted from the Environmental Protection Agency's (EPA) 2011 and 2018 modeling platform (version 6), for GLGE (Source Code Classifications 2260004021–2265004071), and equipment population data were obtained from the EPA's nonroad model. Data were sorted by equipment type and characteristics. Aggregate and equipment-specific emissions were calculated and compared with emissions from all gasoline-fueled nonroad equipment. Results are presented as descriptive statistics. **Results:** In 2011, approximately 26.7 million tons of pollutants were emitted by GLGE (VOC=461,800; CO=5,793,200; NO_x=68,500, PM₁₀=20,700; CO₂=20,382,400), accounting for 24%–45% of all nonroad gasoline emissions. Gasoline-powered landscape maintenance equipment (GLME; leaf blowers/vacuums, and trimmers, edgers, brush cutters) accounted for 43% of VOCs and around 50% of fine PM. Two-stroke engines were responsible for the vast majority of fine PM from GLME. State data (California, New York, Texas, Illinois, and Florida), 2018 projections, and additional comparisons are presented. Methodological issues are discussed. **Conclusions:** GLGE accounts for a major portion of US nonroad gasoline emissions. Two-stroke engines are an important source of VOCs and criteria pollutants.

* Corresponding Author: jamie@quietcommunities.org

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER



INTRODUCTION

Gasoline-powered lawn and garden equipment (GLGE) ranging from string trimmers to stump grinders and tractors is a source of high levels of localized emissions that includes hazardous air pollutants, criteria pollutants, and carbon dioxide (CO₂) [1-4]. Workers using commercial equipment are exposed when they are close to the emitting sources several hours each day, several days a week in seasons of use. Other members of the public, including children, may also be exposed to high levels of emissions from commercial landscape maintenance equipment (GLME) such as leaf blowers, trimmers, and mowers, used routinely around residential neighborhoods, schools, parks, and other public spaces. The commercial landscape maintenance industry has experienced strong growth over the last 15 years and depends largely on gasoline-powered equipment for most tasks once performed manually. These factors are raising concerns about the health impacts of GLGE emissions on workers and the public.

Extensive evidence exists on the adverse health effects of exhaust emissions and other fine particulates which include cardiovascular disease, stroke, respiratory disease, cancer, neurological conditions, premature death, and effects on prenatal development [5-13]. Short term and long term exposures are implicated. However, GLGE as a source of these emissions has received little attention.

Understanding the characteristics of GLGE and GLME emissions can help estimate potential health impacts of these close-to-the-source emissions.

The goal of this study was to characterize annual emissions from GLGE at the national level and in selected states and to estimate the contribution of GLME to those emissions. Special attention is paid to 2-stroke GLME engines. The emissions contributions from the four of the five most populated states are derived from the NEI, and for California, from the emissions inventory of the California Air Resources Board (CARB).

METHODS

Study Design

The GLGE emissions analyzed are total volatile organic compounds (VOC) and individual VOCs (benzene, 1,3 butadiene, acetaldehyde, formaldehyde); criteria pollutants (carbon monoxide [CO], nitrogen oxides [NO_x], particulate matter [PM] <10 microns, including PM < 2.5 microns [PM 10, PM2.5]); and carbon dioxide (CO₂). Equipment pollutant data were extracted from SCC summary reports from the EPA's 2011 and 2018 modeling platform (version 6), and equipment population data were obtained from the Nonroad model. GLGE included the equipment in **TABLE 1** and identified by Source Code Classifications 2260004021–2265004071. The GLME subset is defined as leaf blowers/vacuums; trimmers/edgers/brush cutters; and mowers. Groupings of equipment, eg, leaf blowers/vacuums, were predefined by the NEI.

“All Emissions” are defined as all emissions from stationary and mobile sources, excluding biogenic and naturally occurring emissions. “All Nonroad Emissions” are defined as all emissions from the equipment types accounted for within the Nonroad model; note that this does not include emissions from commercial marine, rail, and aircraft sources. “Gasoline Nonroad Emissions” are defined as emissions from gasoline fueled equipment accounted for within the Nonroad model. National

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER



emissions were analyzed by type of equipment and engine configuration as shown in **TABLE 1**. All results are presented as descriptive statistics.

TABLE 1. Categorization Scheme For Analysis Of GLGE Emission

TYPE OF EQUIPMENT	ENGINE CONFIGURATION
GLME	
Leaf Blowers/Vacuums	2 stroke, 4 stroke
Trimmers/Edgers/Cutters	2 stroke, 4 stroke
Mowers	4 stroke
Other GLGE	
Chain Saws	2 stroke, 4 stroke
Rotary Tillers	2 stroke, 4 stroke
Snowblowers	2 stroke, 4 stroke
Turf Equipment	2 stroke, 4 stroke
Chippers/stump grinders	4 stroke
Tractors	4 stroke
Shredders	4 stroke
Other	4 stroke

Analyses

All analyses except for the 2018 projections represent 2011 estimates.

Equipment Populations

The national populations of all types of GLGE were obtained from the Nonroad model. The contribution of each type to the whole population was determined.

Contributions of All Nonroad and GLGE Sources

All Nonroad Emissions were compared to All Emissions. GLGE emissions were then calculated and compared with All Nonroad Emissions and All Emissions.

Contribution of Landscape Maintenance Equipment to GLGE Emissions

GLME emissions and their contribution to GLGE and All Nonroad Emissions were analyzed. Additional analyses were conducted to examine the relative contributions of 2-stroke GLME engine emissions.

Projected Growth of GLGE Emissions: 2011–2018

GLGE emissions projected for 2018 were obtained from the EPA’s 2018 modeling platform, version 6, and compared with 2011 emissions.



GLGE Emissions in the Five Largest States

State level emissions data from the five most populated states (US Census) – California, Florida, Illinois, New York, and Texas – were extracted and analyzed. Estimates of GLGE emissions for Florida, Illinois, New York, and Texas were based on 2011 data from the EPA’s 2011 modeling platform, version 6. Estimates of GLGE emission for California were based on data from the CARB’s OFFROAD2007 Model and estimated for 2012. No adjustments were made for potential differences in annual emissions between 2011 and 2012 California data. The program structure of the OFFROAD2007 Model provides a general overview of the methodology used to estimate emissions from off-road sources (http://www.arb.ca.gov/msei/offroad/pubs/offroad_overview.pdf).

Each state’s contribution to national GLGE Emissions was calculated and compared with its contributions to the US landscape maintenance labor force and the US population. Labor force statistics were sourced from the Bureau of Labor Statistics, May 2013 reports (www.bls.oes) and population data from the 2011 US Census.

Nonroad Air Emissions Model

EPA developed a nonroad air emissions model in the 1990s to provide estimates of emissions from most types of nonroad equipment, including construction equipment, recreational marine vessels, and lawn and garden equipment (LGE). The model is referred to simply as the “Nonroad” model, and it has been updated a number of times since its creation. Documentation for the model exists as a number of technical reports available on EPA’s website (<http://www.epa.gov/otag/nonrdmdl.htm>). Total emissions are determined by summing the exhaust and evaporative emission components.^{14, 15} The preponderance of emissions from Nonroad equipment occurs as exhaust emissions due to the combustion of fuel. The methodologies for determining exhaust emissions are summarized below.

Exhaust Emissions from Nonroad Engines

The Nonroad model uses the following equation to calculate exhaust emissions from nonroad engines (ref: Median):

- Emissions = (Pop) x (Power) x (LF) x (A) x (EF)
- Where Pop = Engine population
- Power = Average Power (hp)
- LF = Load factor (fraction of available power)
- A = Activity (hrs/yr)
- EF = Emission factor (g/hp-hr)

The derivation of the default model data for each factor from the above equation is discussed below.

a. Equipment Populations And Average Power (Horsepower)

The technical report titled “Nonroad Engine Population Estimates” [16] indicates that equipment population data for most types of equipment were obtained from Power Systems Research, an independent marketing research firm, although in some instances other data source were used. Of interest for this analysis, for many LGE categories EPA used sales data obtained from equipment

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER



manufacturers during the development of its Phase 1 emission standards for small (less than 25 hp) gasoline fueled nonroad engines. This was done for the following LGE categories: lawn mowers, trimmers/edgers/brush cutters, leaf blowers/vacuums, and chainsaws. The report notes that an equipment population base year of either 1996 or 1998 was used for the LGE types.

Once estimates of equipment populations were derived, information obtained by the state of California was used to divide the equipment between the residential and commercial sectors. This step was needed because of the large difference in usage patterns between these two sectors.

TABLE 2 below contains an extract of data from Table 3 of the Nonroad Engine Population report mentioned above, and illustrates how the split between residential and commercial equipment was apportioned for a number of LGE types.

TABLE 2. Percentage Split Between Residential And Commercial Equipment

SCC CODE	APPLICATION	HORSEPOWER CATEGORIES	RESIDENTIAL (% of equipment population)	COMMERCIAL (% of equipment population)
22xx004010 22xx004011	Lawn mowers	All	96.3	3.7
22xx004025 22xx004026	Trimmers / edgers / cutters	0-1 hp	100	0
		1-3 hp	85.3	14.7
		> 3 hp	0	100
22xx004020 22xx004021	Chainsaws	0-1 hp	100	0
		1-3 hp	97.0	3
		> 3 hp	0	100
22xx004030 22xx004031	Leaf blowers / vacuums	0-1 hp	100	0
		1-3 hp	92.5	7.5
		> 3 hp	0	100

i Geographic allocation of residential LGE Populations (except snowblowers)

The Nonroad model uses US Census data on one and two unit housing to allocate national equipment populations to the county level. The population documentation report mentioned above notes that other variables are likely to also affect the distribution of LGE population, such as average yard size. However, no consistent, reliable data surrogates could be found to apportion the national level equipment populations based on these alternative factors, and so the model relies solely upon US Census data on one and two unit housing to allocate national LGE population data to the county level.



ii *Geographic allocation of commercial L&G Equipment Populations (except snowblowers)*

The Nonroad model uses the number of employees in the landscaping services industry to disaggregate national level LGE population data to the county level. This was accomplished using data from the North American Industry Classification System (NAICS); specifically, for NAICS code 561730, landscaping services.

iii *Equipment population projections*

The Nonroad model enables the user to obtain estimates of emissions for years other than the base year used for equipment populations. This is accomplished by the development of processes to handle the growth in equipment populations due to the purchase of new equipment as years pass, and adjustments made to account for the scrappage of old equipment. The reader is referred to the EPA technical reports “Nonroad Engine Growth Estimates,” [17] and “Calculation of Age Distributions in the Nonroad Model – Growth and Scrappage” [18] for further information on these topics. Both of these reports are available on the EPA website (<http://www.epa.gov/otaq/nonrdmdl.htm>).

b. Activity Levels And Load Factors.

Equipment populations and horsepower levels alone are not sufficient for determining emissions from nonroad equipment; assumptions about frequency and patterns of use must also be made. The EPA report, “Median Life, Annual Activity, and Load Factor Values for Nonroad Engine Emissions Modeling” [19] describes how the Nonroad model assigns default activity levels, in hours per year, and load factors in performing its calculations. Load factors are needed to account for the fact that equipment is not typically used at full power 100% of the time; load factors reflect that and are presented in terms of average percent of full power for the equipment as it is used. The activity levels and load factors for small (< or = to 25 hp) spark-ignition engines for many LGE types was taken from data supplied to EPA during the comment period for the regulation of these engines. **TABLE 3** below contains an extract of the default activity data, in annual hours of equipment use, and load factor data, expressed as fraction of full power, taken from Table 6 of the above mentioned report.

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER



TABLE 3. Example Default Activity Levels And Load Factors For LGE

Equipment type	Use	Activity level (Annual hours)	Load factor (fraction of full power)
Lawn mowers	Residential	25	0.33
	Commercial	406	0.33
Trimmers/Edgers/Cutters	Residential	9	0.91
	Commercial	137	0.91
Leaf blowers\Vacuums	Residential	10	0.94
	Commercial	282	0.94
Chainsaws	Residential	13	0.70
	Commercial	303	0.70

c. Emission Factors

EPA’s documentation for the source of the emission factors used within the Nonroad model are contained in the following two reports: “Exhaust and Crankcase Emission Factors for Nonroad Engine Modeling: Compression-Ignition”²⁰ and “Exhaust Emission Factors for Nonroad Engine Modeling: Spark-Ignition.”²¹ Information pertaining to LGE contained in the latter report is discussed below.

Emission factors for spark-ignition engines rated at less than 25 hp were segregated into 5 engine classes based on primary use of the engine (handheld vs. non-handheld), and engine size according to engine displacement. Beginning in 1997, engines designed for both handheld and non-handheld applications became subject to several phases of regulation geared towards reducing fuel consumption (expressed in terms of brake-specific fuel consumption [BSFC]) and producing fewer air emissions in the combustion process. **TABLE 4** below contains an extract of information from Table 1 of the Exhaust Emissions 2010 report, and shows the impact of EPA’s regulation on one such class of engines: small, hand-held, gasoline fueled two-stroke engines.

TABLE 4: Impact Of Regulation On Small*, Hand-Held, Gasoline Fueled Two Stroke Engines

Engine Tech Type	HC (g/hp-hr)	CO (g/hp-hr)	NOx (g/hp-hr)	PM (g/hp-hr)	BSFC (lb/hp-hr)
Baseline	261.00	718.87	0.97	7.7	1.365
Phase 1	219.99	480.31	0.78	7.7	1.184
Phase 2 (with catalyst)	26.87	141.69	1.49	7.7	0.822



BSFC: Brake-specific fuel consumption; CO: carbon monoxide; HC: hydrocarbon; NOx: nitrogen oxides; PM: particulate matter


- These emission factors are for engines sized from 0 to 1 hp.

Other factors also influence the combustion related exhaust emissions from nonroad engines, such as fuel type, ambient temperature, and deterioration of equipment with age and use. The reader is referred to the EPA web-site (<http://www.epa.gov/otaq/nonrdmdl.htm>) for additional information on these topics.

RESULTS

Equipment Populations

Approximately 121 million pieces of GLGE are estimated to be in use in the United States (FIGURE 1). GLME accounts for two-thirds of all GLGE of which lawn mowers are the most numerous, followed by trimmers/edgers/ brush cutters, and then leaf blowers/vacuums. Projections from 2011 indicate a 13% increase across all equipment types after the combined effect of new equipment purchases and scrappage of old equipment are evaluated, resulting in an estimated 136 million pieces of GLGE in use by 2018.



Contribution of Nonroad Emissions to All Emissions

All Nonroad sources account for approximately 242 million tons of pollutants each year, accounting for 17% of all VOC emissions, 12% of NOx emissions, 29% of CO emissions, 4% of CO2 emissions, 2% of PM10 emissions, and 5% of PM2.5 emissions.




All Nonroad Emissions account for a substantial percentage of All Emissions of benzene (25%), 1,3 butadiene (22%), CO (29%), PM10 (2%), and PM2.5 (5%). Because of the relatively small contribution of GLGE CO2 to All Emissions (0.3%), it is not further considered in this report.

Contribution of GLGE to All Emissions and Nonroad Emissions


GLGE emitted approximately 6.3 million tons of VOCs (461,800) and criteria pollutants (CO=5,793,200; NOx=68,500, PM10=20,700 [19,000 of which is PM2.5]), and 20.4 million tons of CO2 in 2011. GLGE represented nearly 4% of All Emissions of VOCs and 12% of All Emissions of CO

(FIGURE 2). GLGE fine PM emissions constitute a fraction of a percent of All Emissions of fine PM, but is a major Nonroad source, accounting for nearly 13% of All Nonroad Emissions of fine PM and more than one-third of Gasoline Nonroad Emissions of fine PM.




Analysis of individual VOC emissions shows that GLGE contributes nearly 8% of All Emissions of both benzene and 1,3 butadiene (**FIGURE 3**). Within All Nonroad Emissions and Gasoline Nonroad Emissions, GLGE accounts for nearly one-third or more of benzene and 1,3 butadiene emissions, and also becomes a major source of aldehyde and formaldehyde emissions from Gasoline Nonroad sources.






Contribution of GLME to GLGE Emissions

Compared with the GLGE contributions of Nonroad Gasoline Emissions shown in **FIGURE 2**, contributions of VOCs and fine PM emissions from GLME are disproportionately high, and for NOx and CO, are disproportionately low (**FIGURE 4**). Small GLME engines account for more than 40% of VOC emissions and one-half of PM10 and PM2.5 emissions from GLGE. Close to 90% of fine PM emissions from GLME come from 2-stroke engines (**FIGURE 5**).





Projected Growth of GLGE Emissions: 2011–2018

By 2018, the annual tonnage of ozone precursors, VOCs and NOx, emitted by GLGE is projected to decrease substantially from 2011, as more of the in-use fleet becomes represented by equipment built to meet EPA nonroad emission standards. CO emissions remain comparable to 2011 levels, while CO2 and fine PM emissions are projected to increase modestly.


TABLE 5: Estimated Change In GLGE Emissions, 2018 Vs 2011

Emissions	% Change
VOCs	-20.9%
NOx	-31.1%
CO	-4.9%
CO2	12.3%
PM 10	8.2%
PM 2.5	8.4%

GLGE Emissions in the Five Most Populated States

When considered together, GLGE emissions from California, Florida, Illinois, New York and Texas constitute approximately one-quarter of national GLGE emissions.






Florida’s GLGE emissions were 1.4 to 2.1-times higher compared with emissions in states having the next highest level of emissions in each GLGE pollutant category, and 2.2 to 4.4-times higher compared with emissions in states having the lowest level of emissions in each GLGE pollutant category (**FIGURE 6**).

For Florida, Illinois, and New York, state-specific contributions of GLGE emissions compared to the national total were relatively consistent with their contributions to the national population and the national grounds maintenance workforce. For California, its GLGE emission contribution was one-fifth that of its contribution to the national population and to the national grounds maintenance workforce.

For Texas, its GLGE emission contribution was 40%–50% that of its contribution to the national population and to the national grounds maintenance workforce (**FIGURE 7**).

- INDEX
- OVERVIEW
- FACTS
- QTC
- ZONES
- PROJECTS
- TEAM
- SLIDES
- REPORT (17p)
- AD 1
- AD 2
- FLYER





DISCUSSION

The main findings of this study are: 1) GLGE is a prevalent source of toxic and carcinogenic emissions; 2) GLGE contributes substantially to nonroad emissions of benzene, 1,3 butadiene, formaldehyde, CO, and fine PM; 3) GLME accounts for a disproportionately large share of VOC and fine PM emissions; 4) 2-stroke engines account for most fine PM emissions from GLME; 5) VOCs and NOx are projected to decrease substantially by 2018; CO emissions remain comparable to 2011 levels; and CO2 and fine PM emissions are projected to increase modestly; and 6) the GLGE emissions contributions from the largest states are not always consistent with contributions to national population and national grounds maintenance workforce.

The large volume of emissions from GLGE found in this study is consistent with findings previously reported by the EPA [1] and from other studies [2-4]. The very substantial contribution of VOC, in particular benzene and 1,3 butadiene, deserves attention especially because of their localized nature.

While VOC emissions are expected decrease 21% on average by 2018, the rates of equipment replacement on which those projections are based are only approximated.

Adverse health effects from the GLGE emissions are well known. Benzene, 1,3 butadiene, and formaldehyde are listed among the four top ranking cancer-causing compounds [22]. They cause lymphomas, leukemias, and other types of cancer (International Agency for Research on Cancer, World Health Organization) [23, 24]. Ground level ozone (formed by VOCs and NOx in the presence of sunlight) and fine PM cause or contribute to early death, heart attack, stroke, congestive heart failure, asthma, chronic obstructive pulmonary disease, and cancer [5-11]. Growing evidence suggests these pollutants also contribute to developmental and neurological

- INDEX
- OVERVIEW
- FACTS
- QTC
- ZONES
- PROJECTS
- TEAM
- SLIDES
- REPORT (17p)
- AD 1
- AD 2
- FLYER



disorders, including autism [7-9, 12, 13]. The mounting evidence on the dangers of short term exposure are especially concerning [7, 9, 11].

The high levels of VOCs and fine PM from GLME are health risks for workers and other members of the public close to the emitting source. Although no studies of grounds maintenance workers were found, studies of gas station workers have shown that regular exposure to gasoline vapors can produce hematological and immunological abnormalities and elevate the risk of cancer [25-27]. In addition, children, seniors, and persons with chronic illnesses are especially vulnerable to the negative health impacts of GLME emissions [28]. Routine use of GLME in the vicinity of residential neighborhoods, schools, parks, and other public spaces may be exposing the public to unnecessary and preventable health risks. New equipment standards do not affect fine PM emissions; in fact, those emissions are expected to increase.

School buses represent another example of a close-to-emitting source in which children are subjected to increased exposure from diesel exhaust [29]. Tests of school buses found that diesel exhaust entering through the front door of the bus results in elevated levels of PM over time. When queuing, PM built up rapidly in the bus cabin when the front doors were open.

The variation in emissions levels observed among the five most populated states should be explored further. The reasons for the high emissions contribution from Florida and relatively low emissions contributions from Texas and California are not clear. Differences between CARB data and NEI data may account for some of the difference between California and other states. For example, the NEI baseline equipment population data are older compared with those of CARB. Other factors that may be involved include but are not limited to emissions estimation procedure, geographic and climate factors, regulations and their effectiveness, and efforts to promote cleaner alternatives.

This study has several limitations. Not all potentially harmful emissions were characterized; for example, polycyclic aromatic hydrocarbons. Other limitations concern the source data. Although the NEI is a comprehensive source of GLGE emissions data, the accuracy of the reported data is uncertain. Baseline equipment population data for the Nonroad model is 15–20 years old and does not account for growth of the commercial industry. This older population data supplies emission estimates to NEI, which in turn is used to create EPA’s 2011 and 2018 modeling platforms. Although the residential and commercial CARB inventories and activity data are newer, they depend largely upon telephone survey data [30, 31]. Methodological weaknesses with the commercial survey data are discussed in the survey report [31]. For both data sources, the rates of replacement of older equipment by newer, cleaner equipment that meets the newer Phase 3 standards [32] can only be approximated.

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER



CONCLUSIONS

GLGE is an important source of toxic and carcinogenic exhaust and fine particulate matter. Improved reporting and monitoring of localized GLGE emissions should be implemented. Medical and scientific organizations should increase public awareness of GLGE and GLME and identify GLGE as an important local source of dangerous air pollutants. Communities and environmental, public health, and other government agencies should create policies and programs to protect the public from GLGE air pollutants and promote non-polluting alternatives.

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER



REFERENCES

1. Michaels H. "NONROAD Overview," Presented at the 2012 International Emission Inventory Conference of the US Environmental Protection Agency, Tampa, Florida, August 16, 2012.
2. Volckens J, Braddock J, Snow RF, et al. "Emissions Profile From New and In-Use Handheld, 2- Stroke Engines," Atmospheric Environment 2007;41:640-649.
3. Volckens J, Olson DA, Hays MD. "Carbonaceous Species Emitted from Handheld Two-Stroke Engines," Atmospheric Environment 2008;42:1239-1248.
4. Shipchandler R. VOC Emissions from Gas Powered Leaf Blowers in the Chicago Metropolitan Region. Waste Management and Research Center Report. Illinois Waste Management and Research Center, February 2008, TN08-093.
5. American Heart Association. Facts: Danger in the Air -Air Pollution and Cardiovascular Disease. Accessed 1/6/14 at http://www.heart.org/HEARTORG/Advocate/IssuesandCampaigns/Advocacy-Fact-Sheets_UCM_450256_Article.jsp
6. American Lung Association. State of the Air 2014.
7. *Integrated Science Assessment for Particulate Matter- Final Report*, US Environmental Protection Agency, December 2009, EPA/600/R-08/139F.
8. Provisional Assessment of Recent Studies on Health Effects of Particulate Matter Exposure, US Environmental Protection Agency, December 2012, EPA/600/R-12/056F,.
9. Integrated Science Assessment for Ozone and Related Photochemical Oxidants, US Environmental Protection Agency, 2013, EPA/600/R-10/076F.
10. *Air Pollution and Cancer*, K Straif, A Cohen, J Samet (Eds), Scientific Publication 161, International Agency for Research in Cancer, World Health Organization, Lyon Cedex FR:IARC, 2013.
11. Shah ASV, Lee KK, McAllister DA, et al. "Short Term Exposure to Air Pollution and Stroke: Systematic Review and Meta-Analysis," BMJ 2015;350:h1295.
12. Raz R, Roberts AL, Lyall K, Hart JE, Just AC, Laden F, Weisskopf MG. "Autism Spectrum Disorder and Particulate Matter Air Pollution Before, During, and After Pregnancy: A Nested Case- Control Analysis within the Nurses' Health Study II Cohort," Environ Health Perspect. 2015 Mar;123(3):264-70.
13. Power MC, Kioumourtoglou M-A, Hart JE, et al. "The Relation Between Past Exposure to Fine Particulate Pollution and Prevalent Anxiety: Observational Cohort Study," BMJ 2015;350:h1111.
14. *Nonroad Evaporative Emission Rates*, US Environmental Protection Agency, July 2010, EPA-420- R-10-021, NR-012-d.
15. *Refueling Emissions for Nonroad Engine Modeling*, US Environmental Protection Agency, April 2004, EPA420-P-04-013, NR-013-b.
16. *Nonroad Engine Population Estimates*, US Environmental Protection Agency, July 2010, EPA-420- R-10-017, NR-006e.
17. *Nonroad Engine Growth Estimate*, US Environmental Protection Agency, April 2004, EPA420-P- 04-008, NR-008c.
18. *Calculation of Age Distributions in the Nonroad Model: Growth and Scrappage*, US Environmental Protection Agency, December 2005, EPA420-R-05-018, NR-007c.
19. Median Life, Annual Activity, and Load Factor Values for Nonroad Engine Emissions Modeling, US Environmental Protection Agency, July 2010, EPA-420-R-10-016, NR-005d.



20. Exhaust and Crankcase Emission Factors for Nonroad Engine Modeling: Compression-Ignition, US Environmental Protection Agency, July 2010, EPA-420-R-10-018, NR-009d.
21. *Exhaust Emission Factors for Nonroad Engine Modeling: Spark-Ignition*, US Environmental Protection Agency, July 2010, EPA-420-R-10-019, NR-010-f.
22. Loh MM, Levy JI, Spengler JD, et al. “Ranking Cancer Risks of Organic Hazardous Air Pollutants in the United States,” *Environ Health Perspect* 2007; 115:1160–1168.
23. Baan R, Gross Y, Straif K et al on behalf of the WHO International Agency for Research on Cancer Monograph Working Group. “A Review of Human Carcinogens—Part F: Chemical Agents and Related Occupations,” *Lancet Oncology* 2009; 10:1143-1144.
24. Report on Carcinogens 13th Edition, 2014. US Department of Health and Human Services, Public Health Service, National Toxicology Program.
25. Lynge E, Andersen A, Nilsson R, et al. “Risk of Cancer and Exposure to Gasoline Vapors,” *Am J Epidemiol* 1997;145:445-458.
26. Tunsaringkarn T, Prueksasit T, Kitwattanavong M, et al. “Cancer Risk Analysis of Benzene, Formaldehyde and Acetaldehyde on Gasoline Station Workers,” *Journal of Environmental Engineering and Ecological Science* 2012, 1:1.<http://dx.doi.org/10.7243/2050-1323-1-1>.
27. Moro AM, Brucker N, Charão MF, et al. “Early Hematological and Immunological Alterations in Gasoline Station Attendants Exposed to Benzene,” *Environ Res.* 2015;137C:349-356.
28. *State of the Air*, 2014, American Lung Association.
29. Hill LB, Zimmerman NJ, Gooch J. A Multi-City Investigation of the Effectiveness of Retrofit Emissions Controls in Reducing Exposures to Particulate Matter in School Buses, Clean Air Task Force, January 2005.
30. 2012 California Survey of Residential Lawn and Garden Equipment Owners: Population and Activity, California Air Resources Board.
31. Acquisition and Analysis of Commercial and Institutional Lawn and Garden Population and Activity Data: Final Report, August 8, 2006, Eastern Research Group, Inc. for the California Air Resources Board.
32. *EPA Finalizes Emission Standards for New Nonroad Spark-Ignition Engines, Equipment, and Vessels*, US Environmental Protection Agency, Office of Transportation and Air Quality, September 2008, EPA420-F-08-013.

INDEX

OVERVIEW

FACTS

QTC

ZONES

PROJECTS

TEAM

SLIDES

REPORT (17p)

AD 1

AD 2

FLYER

